

AD-A091 733

UNION COLL AND UNIV SCHENECTADY NY INST OF ADMINISTRATION--ETC F/6 12/1  
SOME SIMPLE EXAMPLES OF TIME SERIES IN M DIMENSIONS: AN INTRODUCTION--ETC(U)  
SEP 80 L A AROIAN, V TANEJA N00014-77-C-0438

UNCLASSIFIED

AES-8005

NL

1 of 1  
80-1111



END  
DATE  
FILMED  
81-1-1  
DTIC

# LEVEL

125

AD A091733

## SOME SIMPLE EXAMPLES OF TIME SERIES IN M DIMENSIONS:

### AN INTRODUCTION

Leo A. Aroian  
Union College and University  
Institute of Administration  
and Management  
Schenectady, N.Y. 12308

Vidya Taneja  
Western Illinois University  
Professor of Mathematics  
Macomb, Illinois 61455

DEC 6 1980  
NOV 6 1980  
C

#### ABSTRACT

Some simple theoretical examples of time series in  $m$  dimensions, their main properties, estimation, and forecasting are given. These are generalizations of the usual time series to events in both time and space. How these models may be simulated is explained. One practical example of flooding on the Mohawk River is given. Included are new results in forecasting; the regions for permissible values of the autocorrelation and cross correlations; and methods for obtaining these correlations.

#### 1. INTRODUCTION

The purpose of this paper is to give a short introduction to the main ideas of time series in  $m$  dimensions. These simple examples are concerned with results in only one dimension. Full generality to  $m$  dimensions is given in the papers of Aroian and his collaborators. The usual time series is in zero dimensions. The same methods apply to purely spatial series if time is not a consideration; spatial series are explored in a forthcoming paper of Aroian and Gebizlioglu (1). Every event occurs in time and space. An event has many characteristics; only the univariate case is treated here. For the multivariate case, see Aroian (2). As examples, if  $m = 1$ , note the characteristics of a river, and as a generalization any process involving a flow. If  $m = 2$ , examples are physical processes as a storm, rainfall over a region, the changing physical or social processes over time in a restricted region; for  $m = 3$ , examples are the positions of satellites, sunspots on the sun, geological processes and earthquakes, storms in space, or characteristics over an  $(\xi_1, \xi_2, \xi_3, t)$  coordinate system. Social, industrial, scientific, geophysical as well as other processes may be explained by these models.

#### 2. ARMA MODEL

The autoregressive moving average model,  $m = 1$ , is defined for  $m = 1$ ,  $r_1 = r_2 = 2$  by:

$$z_{x,t} = \phi_1 z_{x,t-1} + \phi_2 z_{x-1,t-1} - \theta_1 a_{x,t-1} - \theta_2 a_{x-1,t-1} + a_{x,t} \quad (2.1)$$

The characteristic  $z_{x,t}$  in which one is interested is given as a linear combination of two past values of  $z$  at  $(x,t-1)$  and  $(x-1,t-1)$ ,  $r_2 = 2$ ; and two past values of a variable  $a_{x,t-1}$ ,  $a_{x-1,t-1}$ , random shocks, with an error  $a_{x,t}$ ,  $r_1 = 2$ . We assume  $z_{x,t}$  is weakly stationary,  $E(z_{x,t}) = 0$  (with no loss generality),  $E(z_{x,t}^2) = \sigma_z^2 < \infty$ ,  $E(z_{x,t_1} z_{y,t_2}) = \sigma_z^2 \delta_{x-y, t_1-t_2}$ ,  $-\infty < x < \infty$ ,  $-\infty < t < \infty$ ;  $a_{x,t}$  is an independent random variable with mean zero, variance  $E(z_{x,t_1} z_{x,t_2}) = E(a_{x,t_1} a_{x,t_2}) = 0$ , unless  $t_1 = t_2 = 0$ , when it is  $\sigma_a^2$ ; and  $a_{x,t}$  is independent of  $z_{x-1,t-k}$  unless  $t = k = 0$ .

If  $\phi_1 = \phi_2 = 0$ , then

$$z_{x,t} = a_{x,t} - \theta_1 a_{x,t-1} - \theta_2 a_{x-1,t-1} \quad (2.2)$$

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

DDC FILE COPY

00 11 71 126

# ERRATA SHEET

## SOME SIMPLE EXAMPLES OF TIME SERIES IN M DIMENSIONS: AN INTRODUCTION

by

Leo A. Aroian and Vidya Taneja

- P. 1, l4†  $Ea_{x,t}^2 = \sigma_a^2$
- P. 2, l2† (2.3)  $z_{x,t} = \phi_1 z_{x,t-1} + \phi_2 z_{x-1,t-1} + a_{x,t}$
- P. 2, l3†  $\theta_2 = -\rho_{10}/\rho_{01}$
- P. 3 (4.1)  $z_{x,t} = a_{x,t} - \theta_1 a_{x,t-1} - \theta_2 a_{x-1,t-1}$
- P. 3, l4†  $r = s = 10$
- P. 4, l22† Replace (3) by (2)
- P. 4 (5.1)  $\hat{\rho}\hat{\theta}_1, \hat{\theta}_2 = -r_{10}$
- P. 6 (6.7)  $\phi_1 = (\rho_{01} - \rho_{11}\rho_{10})(1 - \rho_{10}^2)^{-1}$
- P. 6, l3† (6.5) and (6.9) are satisfied.
- P. 7 (6.10)  $\hat{\rho}\hat{\phi}_1, \hat{\phi}_2 = -r_{10}$
- P. 9 (7.6)  $\psi_{ii} = \phi_2 \psi_{i-1,i-1}$
- P. 10 (8.5)  $\sigma_{e_{x,t}}^2 (1,2)$
- P. 10 (8.10)  $\sigma_{e_{x,t}}^2 (l_1, l_2)$
- P. 11 (8.13)  $\sigma_{e_{x,t}}^2 (l_1, l_2)$
- P. 12 Add (10) Campbell, D. & Schaeffer, D.J. (1979). Pollution in the Chicago Sanitary and Ship Canal. Environmental Management, 3, 283-288.

is a moving average MA model  $m_1 = 1$ ,  $r_1 = 2$ ; if  $\theta_1 = \theta_2 = 0$

$$a_{x,t} = \theta_1^2 x_{t-1} + \theta_2^2 x_{t-1,t-1} + a_{x,t} \quad (2.3)$$

is an autoregressive AR model  $m = 1$ ,  $r_2 = 2$ .

If the  $t$  variable is omitted and if in Equation (2.1) one replaces  $x$  and  $x-1$  on the right hand side by  $x-1$  and  $x-2$  except in  $a_{x,t}$  one gets a purely spatial model; while if the  $x$  variable is omitted, one gets the usual time series. A simpler ARMA model is obtained if one sets  $\theta_1 = \theta_2 = 0$ , or  $\theta_1 = \theta_2 = 0$ . The MA model, the AR model, and the ARMA model with their properties will be considered.

### 3. PROPERTIES OF THE MA MODEL

The autocorrelation function of the MA model, the values which  $\theta_1$  and  $\theta_2$  may have, and the corresponding AR model with an infinite set of coefficients will be given. From Voss et al. (3), the variance

$$\sigma_a^2 = \sigma_a^2(1 + \theta_1^2 + \theta_2^2) \quad (3.1)$$

and the correlations

$$\begin{aligned} \rho_{01}(1 + \theta_1^2 + \theta_2^2) &= -\theta_1, \quad \rho_{10}(1 + \theta_1^2 + \theta_2^2) = \theta_1\theta_2, \\ \rho_{11}(1 + \theta_1^2 + \theta_2^2) &= -\theta_2, \end{aligned} \quad (3.2)$$

all other autocorrelations are zero. Note  $\rho_{11} = \rho_{-1-1}$ ,  $\rho_{01} = \rho_{0-1}$ ,  $\rho_{10} = \rho_{-10}$ .

This cutoff property of the autocorrelation function aids in the identification of an MA model  $m = 1$ ,  $r_1 = 2$ . A corresponding cutoff property is true for any MA model,  $(m, r)$ . Rewrite (2.2) as

$$z_{x,t} = (1 - \theta_1 B_t - \theta_2 B_x B_t) a_{x,t} \quad (3.3)$$

where  $B_t a_{x,t} = a_{x,t-1}$ ,  $B_x a_{x,t} = a_{x-1,t}$ , and  $1 - \theta_1 B_t - \theta_2 B_x B_t$  is called the generating function or characteristic polynomial of the MA model. Rewrite (3.3) as

$$a_{x,t} = (1 - \theta_1 B_t - \theta_2 B_x B_t)^{-1} z_{x,t} = \sum_{j=0}^{\infty} (\theta_1 + \theta_2 B_x)^j B_t^j z_{x,t}. \quad (3.4)$$

This is an infinite AR model representation of (2.2) provided  $|\theta_1| + |\theta_2| < 1$ , since  $|B_x| < 1$ ,  $|B_t| < 1$  for convergence of (3.4). Every MA process is stationary but for a representation as an infinite AR model, the condition  $|\theta_1| + |\theta_2| < 1$  is necessary. Note the coefficients of  $z_{x-1,t-k}$  approach zero as  $k \rightarrow \infty$ . These results may also be obtained by successive substitutions in (2.2), or by straightforward division. From (3.2) we infer

$$1 + \theta_1^2 + \theta_2^2 = -\theta_1/\rho_{01} = \theta_1\theta_2/\rho_{10} = -\theta_2/\rho_{11}. \quad (3.5)$$


$$\theta_1 = -\rho_{10}/\rho_{11}, \quad \theta_2 = -\rho_{10}/\rho_{01}, \quad \theta_1 = (\theta_2\rho_{01})/\rho_{11}. \quad (3.6)$$

$$\theta_2 = \frac{-\rho_{11} \pm \rho_{11} [1 - 4(\rho_{01}^2 + \rho_{11}^2)]^{1/2}}{2(\rho_{01}^2 + \rho_{11}^2)} \quad (3.7)$$

$$\text{Hence } 1 - 4(\rho_{01}^2 + \rho_{11}^2) > 0, \text{ or } \rho_{01}^2 + \rho_{11}^2 < 1/4. \quad (3.8)$$

a circle of radius  $1/2$ . From  $\theta_2 = -\rho_{10}/\rho_{01}$ , we conclude  $\rho_{10} = -\rho_{01}\theta_2$ . From  $|\theta_1| + |\theta_2| < 1$ , the added restrictions apply:

$$-1/\rho_{10} < 1/\rho_{11} + 1/\rho_{01} < 1/\rho_{10} \text{ if } 0 < \rho_{10} < 1, \text{ or}$$



$$1/\rho_{10} < 1/\rho_{11} + 1/\rho_{01} < -1/\rho_{10} \text{ if } -1 < \rho_{10} \leq 0. \quad (3.9)$$

Thus the only values of  $\{\rho_{01}, \rho_{10}, \rho_{11}\}$  possible are those in the intersection of (3.8) and (3.9).

In summary, given  $\theta_1$  and  $\theta_2$  then  $\rho_{01}$ ,  $\rho_{10}$ , and  $\rho_{11}$  may be found from (3.2).

If a representation of (2.2) as an infinite AR model (3.4) is desired, then  $|\theta_1| + |\theta_2| < 1$  and the coefficients of this AR model are given by (3.4).

Additionally  $\{\rho_{01}, \rho_{10}, \rho_{11}\}$  satisfy (3.6), (3.7), (3.8), and (3.9).

Conversely, given  $\rho_{01}, \rho_{10}, \rho_{11}$ , which define an MA process (2.2), then if

(3.4) is satisfied, all equations (3.5), (3.6), (3.7), and (3.8) are satisfied. Note  $\rho_{01}$  and  $\rho_{11}$  must satisfy (3.7) and consequently  $\rho_{10} = -\theta_2 \rho_{01}$ . Hence, given  $\{\rho_{01}, \rho_{11}\}$ ,  $\rho_{10}$  is determined. If an arbitrary set

$\{\rho_{01}, \rho_{10}, \rho_{11}\}$  are from some unknown time series in  $m$  dimensions the equations

(3.5), (3.6), (3.7), (3.8), and  $|\theta_1| + |\theta_2| < 1$  will not be satisfied. For

instance, let  $\theta_1 = .2$ ,  $\theta_2 = -.5$ ,  $\sigma_a^2 = 1.29\sigma_a^2$ ,  $\rho_{01} = -.1550$ ,  $\rho_{10} = -.0775$ ,

$\rho_{11} = .3876$ . All equations (3.5-3.9) are satisfied. The corresponding AR

model is  $a_{x,t} = z_{x,t} + .2z_{x,t-1} - .5z_{x,t-2} + .04z_{x,t-3} - .2z_{x,t-4} + .25z_{x,t-5} + .008z_{x,t-6} - .06z_{x,t-7} + .15z_{x,t-8} - .125z_{x,t-9} + \dots$  etc. Suppose from a

simulation of this process the sample values are  $r_{01} = -.14$ ,  $r_{10} = -.10$ ,  $r_{11} = .40$ , to determine the estimated values of the constants  $\hat{\theta}_1, \hat{\theta}_2$  three

equations must be satisfied, an impossibility. Instead iterations are used

in order to satisfy the (3.5) as closely as possible. The result is  $\hat{\theta}_1 = .22$ ,

$\hat{\theta}_2 = -.55$ ,  $1 + \hat{\theta}_1^2 + \hat{\theta}_2^2 = 1.3509$ , and resulting  $r$ 's from (3.5) are  $r_{01} = -.163$ ,

$r_{10} = -.090$ ,  $r_{11} = .407$ .

We rewrite

$$\begin{aligned} a_{x,t} &= \sum_{i,j} \pi_{ij} z_{x-i,t-j}, \quad \pi_{00} = 1, \quad \pi_{01} = \theta_1, \quad \pi_{11} = \theta_2, \\ \pi_{02} &= \theta_1^2, \quad \pi_{12} = 2\theta_1\theta_2, \quad \pi_{22} = \theta_2^2, \quad \pi_{03} = \theta_1^3, \quad \pi_{13} = 3\theta_1^2\theta_2, \\ \pi_{23} &= 3\theta_1\theta_2^2, \quad \pi_{33} = \theta_2^3, \quad \pi_{0j} = \theta_1^j \pi_{0j-1}, \quad \pi_{ij} = \theta_1 \pi_{i,j-1} + \\ &\quad \theta_2 \pi_{i-1,j-1}, \quad \pi_{jj} = \theta_2 \pi_{j-1,j-1}. \end{aligned} \quad (3.10)$$

#### 4. SIMULATION OF THE MA MODEL

Given  $\{\theta_1, \theta_2\}$ ,  $|\theta_1| + |\theta_2| < 1$ , simulate

$$z_{x,t} = a_{x,t} - \theta_1 a_{x,t-1} - \theta_2 a_{x,t-2} \quad (4.1)$$

by first generating a set of random numbers  $\{a_{x+i,t+j}\}$ ,  $i=0,1,2,\dots, r+4, j=0,1,2,\dots, s+4$ , from a distribution  $\mu=0, \sigma=1$ . This gives  $(r+5)(s+5)$  values of  $a_{x+i,t+j}$

random numbers. Then from these values of  $a$ 's calculate  $(r+5)(s+5)$  values of  $\{z_{x+i,t+j}\}$  with the chosen theoretical set  $\{\theta_1, \theta_2\}$  using (4.1). All  $a$ 's not

generated as above are identically zero. Delete the first two rows, first two columns, the last three rows and the last three columns of the  $Z$  matrix.

This produces a reduced  $Z$  matrix of  $rs$  value. Note the number five is chosen for convenience, so an inner  $Z$  matrix may represent the process properly. In

the example  $\theta_1 = .2$ ,  $\theta_2 = -.5$ , and the  $a_{x,t}$ 's are normally distributed with  $\mu=0$ ,

$\sigma_a=1$ ,  $r=s=10$ ,  $z_{x,t}$  is given as the first number in each cell of Table 1. The

$a_{x,t}$ , the second number in each cell of Table 1 is not the  $10 \times 10$  inner part of  $15 \times 15$  original  $Z$  but is determined from the  $10 \times 10$   $Z$  matrix by using

$$a_{x,t} = z_{x,t} + \theta_1 a_{x,t-1} + \theta_2 a_{x,t-2} \quad (4.2)$$

with  $\theta_1 = .2$ ,  $\theta_2 = -.5$  where now  $a_{x,t-1}$  and  $a_{x-1,t-1}$  are taken as zero for initial starting values. Thus in the first row  $a_{x,t} = z_{x,t}$ . In the first column all unknown  $a$ 's for  $x < 1$  are zero. In this way a set of  $a$ 's are built up slightly different from the inner  $10 \times 10$   $a_{x,t}$ 's obtainable from the  $15 \times 15$  original  $Z$  matrix. This is done since in a sample of  $z_{x,t}$  the  $a_{x,t}$  must be determined in this way (although in practice the  $\{\theta_1, \theta_2\}$  would be unknown and would be a set  $\{\hat{\theta}_1, \hat{\theta}_2\}$  estimated from the sample  $z_{x,t}$ 's). In the second row all needed  $a$ 's are known as soon as the first  $a$  is determined, and subsequently for each row, since the needed  $a$ 's in any cell are found by using the  $z_{x,t}$  in that cell and the  $a_{x,t-1}$  in the cell directly above and the  $a_{x-1,t-1}$  in the cell one to the left and above using (4.2). This set of  $a$ 's is used in finding all further  $Z$ 's for all subsequent sets  $\{\theta_1, \theta_2\}$ . From the  $10 \times 10$   $Z$  matrix determine  $r_{m,n}$ ,  $\bar{z}$  (the sample mean),  $\hat{\sigma}_z^2$  (the sample variance of  $z$ ,  $r_{01}$ ,  $r_{10}$ ,  $r_{11}$ ,  $\hat{\theta}_1, \hat{\theta}_2$ , using (3.5-3.7) iteratively as in 3. New estimates  $z_{x,t}$  may be found using any desired  $(\hat{\theta}_1, \hat{\theta}_2)$  particularly a set satisfying (3.5-3.7) and having a minimum prediction error variance  $\hat{\sigma}_e^2 = E(z_{x,t} - \hat{z}_{x,t})^2 / (n-1)$ . This is essentially a least squares procedure. The minimum variance procedure is to vary a set  $(\theta_1, \theta_2)$  in the plane such that  $\hat{\sigma}_e^2$  is a minimum. The theoretical value of  $\sigma_e^2$  is

$$\sigma_e^2 = \sigma_a^2 \{(\theta_1 - \hat{\theta}_1)^2 + (\theta_2 - \hat{\theta}_2)^2\}$$

where  $\theta_1$  and  $\theta_2$  are population values. This method is remarkably effective, working well for autoregressive models also (3). Note  $\hat{z}(x,t)$  are values from the minimum prediction error variance set  $\hat{\theta}_1 = .208, \hat{\theta}_2 = -.493$  almost identical to the theoretical set  $\{.2, -.5\}$ ; while  $\hat{z}(x,t)$  are from  $\{.333, -.418\}$  the least squares set most nearly satisfying (3.5)-(3.7), which leads to a theoretical  $\{\rho_{01}, \rho_{10}, \rho_{11}\}$  set of  $\{-.259, -.108, .325\}$  compared with an actual  $\{r_{10}, r_{01}, r_{11}\}$  of  $\{-.255, -.133, .299\}$  from  $z_{x,t}$ . The values of  $r_{m,n}$  in Table 2 are those from  $z_{x,t}$  (the notation of  $z(x,t)$  in Table 1 is used for convenience.) Further from (5.1) in the next section given  $\{.2, -.5\}$ , the 95% confidence interval for  $\theta_1$  is  $\{.036, .364\}$  and for  $\theta_2$  is  $\{-.664, -.336\}$  and  $\{.333, -.418\}$  cuts these limits.

##### 5. ESTIMATION

The variances and covariances of  $(\hat{\theta}_1, \hat{\theta}_2)$  are approximated by use of the sample estimates  $r_{01}, r_{10}, r_{11}$  in a sample of  $n$  of  $\rho_{01}, \rho_{10}, \rho_{11}$  and the relationship of the variance-covariance matrix of MA models which are the same as those of AR models. From the results of Perry and Aroian (4),

$$\begin{aligned} \hat{\sigma}_{\hat{\theta}_1}^2 &= \hat{\sigma}_{\hat{\theta}_2}^2 = n^{-1} (1 - r_{10}^2)^{-1} (1 - \theta_1^2 - \theta_2^2 - 2\theta_1\theta_2 r_{10}), \text{ and} \\ \hat{\rho}_{\hat{\theta}_1, \hat{\theta}_2} &= r_{10}. \end{aligned} \quad (5.1)$$

These correspond to the least squares solution of the corresponding AR model and if the  $a_{x,t}$  are assumed to be distributed normally, these are close approximations to the maximum likelihood estimates of  $\{\theta_1, \theta_2\}$ . (See Box and Jenkins, (5) p. 283, for similar case,  $m=0$ .) The starting values of  $a_{x,t}$  as noted in the simulation would need to be considered for the maximum likelihood solution. The method leading to (5.1) is general and applies to the  $m$  dimensional MA model. The minimum variance estimates  $\{\hat{\theta}_1, \hat{\theta}_2\}$  and the resulting estimates of the correlation between  $\hat{\theta}_1$  and  $\hat{\theta}_2$  may be found as noted in the next section for any set of data.

TABLE 1  
Values of  $z(x,t)$ ,  $a(x,t)$ ,  $\hat{z}(x,t)_1$ ,  $\hat{z}(x,t)_2$

$t \ x$	1	2	3	4	5	6	7	8	9	10
1 $z(x,t)$	-1.478	0.815	-0.301	0.337	-1.277	0.527	0.210	-0.625	0.867	-1.668
$a(x,t)$	-1.478	0.815	-0.301	0.337	-1.277	0.527	0.210	-0.625	0.867	-1.668
$\hat{z}(x,t)_1$	-1.478	0.815	-0.301	0.337	-1.277	0.527	0.210	-0.625	0.867	-1.668
$\hat{z}(x,t)_2$	-1.478	0.815	-0.301	0.337	-1.277	0.527	0.210	-0.625	0.867	-1.668
2 $z(x,t)$	-1.110	-1.410	1.541	0.126	-0.681	0.245	0.132	1.506	-1.602	-0.741
$a(x,t)$	-1.406	-0.508	1.073	0.343	-1.105	0.989	-0.090	1.276	-1.116	-1.508
$\hat{z}(x,t)_1$	-1.098	-1.406	1.538	0.125	-0.673	0.250	0.127	1.510	-1.605	-0.734
$\hat{z}(x,t)_2$	-0.913	-1.397	1.514	0.106	-0.539	0.280	0.061	1.572	-1.666	-0.590
3 $z(x,t)$	-0.105	-1.199	-0.701	-0.649	1.620	0.425	0.148	-0.124	0.595	-0.060
$a(x,t)$	-0.386	-0.598	-0.232	-1.116	1.228	1.175	-0.364	0.176	-0.266	0.196
$\hat{z}(x,t)_1$	-0.094	-1.185	-0.706	-0.659	1.626	0.425	0.142	-0.134	0.595	-0.040
$\hat{z}(x,t)_2$	0.082	-1.016	-0.802	-0.783	1.739	0.384	0.079	-0.286	0.639	0.232
4 $z(x,t)$	0.450	-0.484	-1.375	-0.278	-1.252	0.540	-0.343	0.767	-2.043	-0.903
$a(x,t)$	0.372	-0.411	-1.123	-0.385	-0.448	0.161	-1.003	0.984	-2.184	-0.731
$\hat{z}(x,t)_1$	0.453	-0.477	-1.369	-0.267	-1.254	0.522	-0.348	0.768	-2.042	-0.903
$\hat{z}(x,t)_2$	0.501	-0.373	-1.295	-0.110	-1.324	0.283	-0.391	0.773	-2.022	-0.907
5 $z(x,t)$	-2.642	0.460	-0.264	-1.163	-0.555	2.482	1.121	-1.062	0.911	-1.187
$a(x,t)$	-2.568	0.191	-0.284	-0.679	-0.452	2.739	0.840	-0.364	-0.018	-0.241
$\hat{z}(x,t)_1$	-2.645	0.461	-0.252	-1.152	-0.549	2.484	1.128	-1.063	0.922	-1.166
$\hat{z}(x,t)_2$	-2.692	0.484	-0.081	-1.020	-0.464	2.497	1.241	-1.111	1.121	-0.911
6 $z(x,t)$	1.355	0.040	-0.401	-0.857	2.023	-1.367	-0.648	0.602	-0.817	-1.329
$a(x,t)$	0.842	1.362	-0.553	-0.851	2.272	-0.593	-1.850	0.109	-0.639	-1.368
$\hat{z}(x,t)_1$	1.376	0.056	-0.400	-0.850	2.031	-1.386	-0.674	0.599	-0.814	-1.327
$\hat{z}(x,t)_2$	1.696	0.225	-0.379	-0.743	2.139	-1.694	-0.984	0.581	-0.785	-1.295
7 $z(x,t)$	0.554	-0.191	1.936	0.196	-0.843	1.234	-0.742	0.557	0.775	0.542
$a(x,t)$	0.722	-0.340	1.144	0.303	0.037	-0.021	-0.815	1.503	0.593	0.588
$\hat{z}(x,t)_1$	0.547	-0.208	1.931	0.207	-0.855	1.223	-0.723	0.569	0.779	0.557
$\hat{z}(x,t)_2$	0.442	-0.441	1.898	0.355	-1.075	1.127	-0.447	0.694	0.851	0.776
8 $z(x,t)$	0.012	2.509	-1.481	0.012	0.692	-0.702	0.124	-2.361	0.689	1.718
$a(x,t)$	0.157	2.080	-1.082	-0.499	0.548	-0.725	-0.029	-1.653	0.056	1.539
$\hat{z}(x,t)_1$	0.006	2.507	-1.488	0.002	0.690	-0.702	0.131	-2.367	0.674	1.709
$\hat{z}(x,t)_2$	-0.084	2.495	-1.605	-0.122	0.662	-0.702	0.234	-2.494	0.487	1.591
9 $z(x,t)$	0.298	-1.145	2.832	0.688	-0.602	-1.427	-1.601	-0.279	-1.305	-0.124
$a(x,t)$	0.329	-0.808	1.576	1.129	-0.243	-1.846	-1.244	-0.596	-0.467	0.156
$\hat{z}(x,t)_1$	0.297	-1.163	2.826	0.700	-0.603	-1.425	-1.596	-0.266	-1.294	-0.137
$\hat{z}(x,t)_2$	0.277	-1.434	2.805	0.843	-0.634	-1.376	-1.538	-0.057	-1.177	-0.333
10 $z(x,t)$	-1.572	1.603	-0.668	1.921	1.776	1.193	-1.140	-0.279	1.135	-1.252
$a(x,t)$	-1.506	1.277	0.051	1.359	1.162	0.945	-0.466	0.224	1.339	-0.988
$\hat{z}(x,t)_1$	-1.575	1.607	-0.675	1.901	1.770	1.209	-1.117	-0.266	1.143	-1.250
$\hat{z}(x,t)_2$	-1.616	1.683	-0.811	1.642	1.716	1.458	-0.823	-0.098	1.246	-1.234

MEAN OF  $z(x,t)$  = -0.0757732      VAR. OF  $z(x,t)$  = 1.29559  
 MEAN OF  $a(x,t)$  = -0.0620963      VAR. OF  $a(x,t)$  = 1.03742  
 MEAN OF  $\hat{z}(x,t)_1$  = -0.0745114      VAR. OF  $\hat{z}(x,t)_1$  = 1.29484

The prediction error variance of  $\hat{z}(x,t)_1$  = .00009444634  
 For  $z(x,t)_1$ ,  $\hat{\theta}_1 = .208$ ,  $\hat{\theta}_2 = -.493$

MEAN OF  $\hat{z}(x,t)_2$  = -0.0575717      VAR. OF  $\hat{z}(x,t)_2$  = 1.32942

The prediction error variance of  $\hat{z}(x,t)_2$  = .0245026  
 For  $z(x,t)_2$ ,  $\hat{\theta}_1 = .333$ ,  $\hat{\theta}_2 = -.418$

Where  $\hat{z}(x,t)_1$  is the prediction with minimum error without regard to (3-5) to (3-8)  
 $z(x,t)_2$  is the prediction with minimum error with regard to (3-5) to (3-8).

TABLE 2  
ESTIMATED CORRELATION COEFFICIENTS  
 $r_{m,n}$  from  $z_{x,t}$

m n	0	1	2	3	4	5
-5	-.02440	-.08929	.01751	.09018	-.07126	.01399
-4	-.00683	-.01955	-.15060	.12190	-.06967	.06822
-3	.00744	-.21941	-.09758	-.16610	.14620	.06586
-2	-.04944	-.02406	.09758	.02137	.03338	.00839
-1	-.13263	.15886	-.05186	.07963	-.08132	.03068
0	1.0000	-.25518	.10146	.01626	-.05956	.02659
1	-.13263	.29921	.03112	-.13549	.03900	-.12985
2	-.04944	.05475	.02902	.05822	-.10431	.04176
3	.00744	-.00829	-.04685	.12587	.01059	.00946
4	-.00683	-.08132	.06358	-.00833	.07500	-.04081
5	-.02440	.00370	-.06670	-.05665	.05308	.01927

#### 6. PROPERTIES OF THE AR MODEL

The properties of the AR model

$$z_{x,t} = \phi_1 z_{x,t-1} + \phi_2 z_{x,t-2} + a_{x,t} \quad (6.1)$$

are taken from Taneja and Aroian (6) and Perry and Aroian (4). These are:  
the characteristic function,

$$\phi(B_x, B_t) = 1 - (\phi_1 + \phi_2 B_x) B_t; \quad (6.2)$$

the autocorrelation function,

$$\rho_{m,n} = \phi_1 \rho_{m,n-1} + \phi_2 \rho_{m,n-2}, \quad m = n \neq 0; \quad (6.3)$$

the variance,

$$\sigma_z^2 = \sigma_a^2 (1 - \phi_1 \rho_{01} - \phi_2 \rho_{11})^{-1}; \quad (6.4)$$

and the stationarity condition

$$|\phi_1| + |\phi_2| < 1; \quad (6.5)$$

the Yule-Walker equations,

$$\rho_{01} = \phi_1 + \phi_2 \rho_{10} \quad \rho_{11} = \phi_1 \rho_{10} + \phi_2 \quad (6.6)$$

with the solution:

$$\begin{aligned} \phi_1 &= (\rho_{01} - \rho_{10} \rho_{10}) (1 - \rho_{10}^2)^{-1} \\ \phi_2 &= (\rho_{11} - \rho_{01} \rho_{10}) (1 - \rho_{10}^2)^{-1} \end{aligned} \quad (6.7)$$

Since  $\sigma_z^2 > 0$ , then  $\{\rho_{01}, \rho_{10}, \rho_{11}\}$  are restricted by

$$\rho_{10}^2 + \rho_{01}^2 + \rho_{11}^2 - 2\rho_{01}\rho_{10}\rho_{11} - 1 < 0. \quad (6.8)$$

If (6.8) is set equal to zero, then for each value of  $\rho_{10}$  (6.8) represents an ellipse. The permissible values lie within the rectangle described in the ellipses. The rectangle is bounded by

$$\begin{aligned} -1 + \rho_{10} < \rho_{01} - \rho_{11} < 1 - \rho_{10}, \text{ and} \\ -1 - \rho_{10} < \rho_{01} + \rho_{11} < 1 + \rho_{10} \end{aligned} \quad (6.9)$$

While an AR model is invertible, it is not stationary unless condition (6.9) is satisfied. Note that given a permissible set of  $\{\rho_{10}, \rho_{11}, \rho_{01}\}$  then  $\phi_1$  and  $\phi_2$  are determined by (6.7). However, if  $(\phi_1, \phi_2)$  are given, then  $\{\rho_{10}, \rho_{11}, \rho_{01}\}$



are unique and may be determined from the corresponding AM model of the AR model. An example, the depths of the Mohawk River, discussed briefly in the Perry and Aroian paper (4) is being expanded from three points on the river to ten points. The results will be used to predict floods along the Mohawk. Let  $\phi_1 = .2$ ,  $\phi_2 = -.6$ ,  $\rho_{10} = -.20871$ ,  $\rho_{01} = .32523$ ,  $\rho_{11} = -.64174$ . The simulation of this set of  $\{\rho_{10}, \rho_{01}, \rho_{11}\}$  is accomplished by using the  $\phi$ 's and any set of starting values with the  $a$ 's generated by any desired distribution. The values of  $\rho_{m,n}$  for this example are given in the table for  $(m,n)$  in the first and second quadrants; the results in the third quadrant are the same as those in the first quadrant, while the results in the fourth quadrant are the same as those in the second quadrant. Thus  $\rho_{m,n} = \rho_{-m,-n}$ ,  $\rho_{m,-n} = \rho_{-m,n}$ ,  $\rho_{m,0} = \rho_{-m,0}$ ,  $\rho_{0,-n} = \rho_{0,n}$ , but  $\rho_{m,n} \neq \rho_{-m,n}$ . This follows from the definition of  $\rho_{x-y, t_1-t_2}$ , given in section 2. A table of the correlation  $\rho_{m,n}$  for  $\phi_1 = .2$ ,  $\phi_2 = -.6$  is presented.

TABLE 3  
CORRELATIONS  $\rho_{m,n}$  IN AR MODEL  
 $\phi_1 = .2$ ,  $\phi_2 = -.6$

n	0	1	2	3	4	5	6	7	8
m									
8								.01	.03
7							-.01	-.04	-.06
6					.01	.02	.07	.09	.07
5				-.01	-.04	-.11	-.12	-.09	-.05
4		.01	.02	.06	.17	.17	.11	.06	.03
3	-.01	-.03	-.09	-.26	-.22	-.12	-.06	-.02	-.01
2	.04	.13	.41	.28	.13	.05	.02	.01	
1	-.21	-.64	-.32	-.13	-.05	-.05	-.02	-.01	
0	1.00	.33	.11	.03	.01				
-1	-.21	-.07	-.02	-.01					
-2	.04	.01							
-3	-.01								

The numbers in the blank spaces are zero.

Every AR model has a cutoff property given by the partial autocorrelation function,  $\phi_i$  in this case;  $\phi_i \neq 0$  for  $i = 1, 2$ ;  $\phi_i = 0$ ,  $i \neq 1, 2$ , where  $\phi_i$  is determined by the Yule-Walker equations corresponding to AR models,  $r > 2$ . For an  $m = 1$  model is of order one if  $\phi_3, \phi_4, \dots, \phi_r$  are zero as determined by the partial coefficient of correlation, Taneja and Aroian (6).

The approximate estimated variances and covariances of  $(\phi_1, \phi_2)$  given by

$$\hat{\sigma}_{\hat{\phi}_1}^2 = \hat{\sigma}_{\hat{\phi}_2}^2 = n^{-1}(1-r_{10}^2)^{-1}(1-\phi_1^2-\phi_2^2-2\phi_1\phi_2 r_{10}) \text{ and} \quad (6.10)$$

$$\hat{\sigma}_{\hat{\phi}_1, \hat{\phi}_2} = r_{10}$$

are approximate maximum likelihood results, if the  $a_{it}$ 's are distributed normally, are asymptotically unbiased, consistent, and minimum variance. Let

$$R_2 = \begin{pmatrix} 1 & r_{10} \\ r_{10} & 1 \end{pmatrix}, \quad r^1 = (r_{01}, r_{11}), \quad \hat{\phi}^1 = (\hat{\phi}_1, \hat{\phi}_2)$$

$$\hat{V}(\hat{\phi}_1, \hat{\phi}_2) = n^{-1}(1-r^2) R_2^{-1} = n^{-1}(1-r_{01}\hat{\phi}_1 - r_{11}\hat{\phi}_2) R_2^{-1},$$

where  $\hat{V}(\hat{\phi}_1, \hat{\phi}_2)$  is the variance-covariance matrix of  $(\hat{\phi}_1, \hat{\phi}_2)$ . The minimum variances of  $(\hat{\phi}_1, \hat{\phi}_2)$  may be found numerically by varying  $(\hat{\phi}_1, \hat{\phi}_2)$  until the variance  $\sigma_a^2$  is minimized for the particular sample.

The AR model may be given as an infinite AM model

$$\begin{aligned} z_{x,t} &= \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \psi_{ij} z_{x-i,t-j}, \\ \psi_{00} &= 1, \psi_{01} = \phi_1, \psi_{11} = \phi_2, \psi_{02} = \phi_1^2, \psi_{12} = 2\phi_1\phi_2, \psi_{22} = \phi_2^2, \\ \psi_{0j} &= \phi_1\psi_{0j-1}, \psi_{ij} = \phi_1\psi_{i,j-1} + \phi_2\psi_{i-1,j-1}, \psi_{jj} = \phi_2\psi_{j-1,j-1}, \end{aligned} \quad (6.11)$$

like (4.1) replacing  $\pi_{ij}$  by  $\psi_{ij}$ , and  $\theta_i$  by  $\phi_i$ .

For  $\phi_1 = .2$ ,  $\phi_2 = -.6$ ,  $\psi_{00} = 1$ ,  $\psi_{01} = .2$ ,  $\psi_{11} = -.6$ ,  $\psi_{02} = .04$ ,

$$\psi_{12} = -.24, \psi_{22} = .36, \psi_{03} = .008, \psi_{13} = -.072, \psi_{23} = .216,$$

$$\psi_{33} = .216, \text{ etc.}$$

An AR model simulation is given in Perry and Aroian (4).

#### 7. ARMA MODELS

The results of Oprian et al. (7) are used freely in addition to some new results mainly in the determination of values of  $(\phi_1, \phi_2, \theta_1, \theta_2)$  as related to the values of  $(\rho_{01}, \rho_{10}, \rho_{11}, \rho_{-1,1})$ . The results are:

$$\begin{aligned} \sigma_z^2 &= \sigma_a^2 \{1 - \theta_1(\phi_1 - \theta_1) - \theta_2(\phi_2 - \theta_2)\} (1 - \phi_1\rho_{01} - \phi_2\rho_{10})^{-1}, \\ \rho_{01} &= \phi_1 + \phi_2\rho_{10} - \theta_1\sigma_a^2/\sigma_z^2; \theta_1\sigma_a^2/\sigma_z^2 = \phi_1 + \phi_2\rho_{10} - \rho_{01} \\ \rho_{11} &= \phi_1\rho_{10} + \phi_2 - \theta_2\sigma_a^2/\sigma_z^2; \theta_2\sigma_a^2/\sigma_z^2 = \phi_1\rho_{10} + \phi_2 - \rho_{11} \\ \rho_{10} &= \phi_1\rho_{1-1} + \phi_2\rho_{01} - \theta_2(\phi_1 - \theta_1)\sigma_a^2/\sigma_z^2. \end{aligned} \quad (7.1)$$

The second and third equations may be solved for  $\theta_1\sigma_a^2/\sigma_z^2$  and these results substituted in the first and fourth equations to obtain  $(\phi_1, \phi_2, \theta_1, \theta_2)$ . If  $\phi_1 = \phi_2 = 0$ , or  $\theta_1 = \theta_2 = 0$ , the corresponding MA or AR models are obtained. For  $m, n \geq 2$ ,

$$\rho_{m,n} = \phi_1\rho_{m,n-1} + \phi_2\rho_{m-1,n-1}. \quad (7.2)$$

The ARMA model may be represented as an infinite moving MA model

$$\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \psi_{ij} z_{x-i,t-j}, \text{ or} \quad (7.3)$$

an infinite AR model

$$\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \pi_{ij} z_{x-i,t-j}, \quad (7.4)$$

restrictions already noted  $|\phi_1| + |\phi_2| < 1$  and  $|\theta_1| + |\theta_2| < 1$ , naturally restricting the values of  $(\rho_{01}, \rho_{10}, \rho_{11})$  also. The infinite MA model is given by

$$z_{x,t} = (1 - \theta_1 B_t - \theta_2 B_x B_t) (1 - \phi_1 B_t - \phi_2 B_x B_t)^{-1} a_{x,t} =$$

$$(1 - \theta_1 B_t - \theta_2 B_x B_t) \left( \sum_{i=0}^{\infty} (\phi_1 + \phi_2 B_x)^i B_t^i \right) a_{x,t}, \quad (7.5)$$

with  $\psi_{00} = 1$ ,  $\psi_{01} = \phi_1 - \theta_1$ ,  $\psi_{11} = \phi_2 - \theta_2$ ,  $\psi_{02} = \phi_1$ ,

$$\psi_{12} = -\phi_1 \phi_2 + 2\phi_1 \phi_2 - \phi_2 \theta_1, \dots, \psi_{0j} = \phi_1 \psi_{0,j-1},$$

$$\psi_{ij} = \phi_1 \psi_{i,j-1} + \phi_2 \psi_{i-1,j-1}, \psi_{ii} = \phi_2 \psi_{i-1,i-1}. \quad (7.6)$$

Similarly the ARMA model as an AR model:

$$\pi_{00} = 1, \pi_{01} = \theta_1 - \phi_1, \pi_{11} = \theta_2 - \phi_2, \pi_{02} = \theta_1 (\theta_1 - \phi_1)$$

$$\pi_{12} = -\theta_1 \phi_2 + 2\theta_1 \phi_2 - \theta_2 \phi_1, \pi_{22} = \theta_2 (\theta_2 - \phi_2), \quad (7.7)$$

$$\pi_{0j} = \theta_1 \pi_{0,j-1}, \pi_{ij} = \theta_1 \pi_{i,j-1} + \theta_2 \pi_{i-1,j-1},$$

$$\pi_{ii} = \theta_2 \pi_{i-1,i-1}.$$

Note (3.10) and (6.11) are special cases of (7.5) and (7.6). For

$$\phi_1 = .2, \phi_2 = -.6, \theta_1 = .2, \theta_2 = -.5,$$

$$\psi_{00} = 1, \psi_{11} = -.1, \psi_{12} = -.02, \psi_{22} = -.6, \psi_{13} = -.004, \quad (7.8)$$

$$\psi_{23} = .024, \psi_{33} = -.036, \text{ etc.}$$

The autocorrelation function must be found by using (7.5) and

$$c_{m,n} = (\sum_{ij} \psi_{ij} \psi_{i-m,j-n}) (\sum_{ij} \psi_{ij}^2)^{-1}, m = n \neq 0 \quad (7.9)$$

and

$$\sigma_z^2 = \sigma_a^2 \sum_{ij} \psi_{ij}^2. \quad (7.10)$$

Thus  $c_z^2 / \sigma_a^2 = 1.01741$ , obtained from (7.9) and (7.10), must be used in (7.1) to solve for  $\{c_{01}, c_{10}, c_{11}, c_{11-1}\} = \{.058, -.091, -.127, -.213\}$  as compared to the MA model  $\{-.155, -.078, .389, 0\}$ , and the AR model  $\{.33, -.21, -.64, -.07\}$ . In fact, had  $\phi_1 = .2, \phi_2 = -.6, \theta_1 = .2, \theta_2 = -.5$ , been chosen, then the ARMA model would have been reduced to  $(1 - .2B_t + .6B_x B_t) z_{x,t} = (1 - .2B_t + .6B_x B_t) a_{x,t}$ , or  $z_{x,t} = a_{x,t}$ . Conversely, had a sample set of  $s_z^2 / s_a^2, \{r_{01}, r_{10}, r_{11}, r_{11-1}\}$  been given, estimates  $(\hat{\phi}_1, \hat{\phi}_2, \hat{\theta}_1, \hat{\theta}_2)$  could be found from (7.1) as already given by substituting the sample estimates for the population values. These results with the exception of (7.1) and the methods of solution given there are due to Robert Perry in his ongoing Ph.D. thesis.

How may the estimates of  $\{\phi_1, \phi_2, \theta_1, \theta_2\}$  in a sample of  $n$  be found? One method is to choose the set  $\{\hat{\phi}_1, \hat{\phi}_2, \hat{\theta}_1, \hat{\theta}_2\}$  which minimizes  $s_e^2$ , the sample estimates of  $\sigma_e^2$  as done in 4 by varying the four constants numerically. The other method is to find the corresponding AR model through  $\pi_{22}$  and then apply the method given in section 5.

## 8. FORECASTING

Now that the models have been defined, their properties given, it is necessary to show how these results may be used for forecasting. It should be realized that in forecasting it is not necessary to assume the future repeats the past. All that is needed is to assume that the errors of the past will be repeated in the future, although the events themselves will not necessarily be

formulas must not be used blindly, but must be checked carefully.

In the MA case (2.1):

$$\hat{z}_{x+l_1, t+l_2} = a_{x+l_1, t+l_2} - \theta_1 a_{x+l_1, t+l_2-1} - \theta_2 a_{x+l_1-1, t+l_2-1}, \quad (8.11)$$

$$e_{x+l_1, t+l_2} = a_{x+l_1, t+l_2} - \theta_1 a_{x+l_1, t+l_2-1} - \theta_2 a_{x+l_1-1, t+l_2-1} \quad (8.12)$$

$$\hat{z}_{x, t}(l_1, l_2) = 0$$

$$\sigma_z^2(l_1, l_2) = \sigma_a^2(1 + \theta_1^2 + \theta_2^2), \quad (8.13)$$

for  $l_1 \geq 2$  or  $l_2 \geq 2$ . The other cases are

$$\hat{z}_{x, t}(0, 1) = -\theta_1 a_{x, t} - \theta_2 a_{x-1, t}, \quad \hat{z}_{x, t}(1, 0) = -\theta_2 a_{x, t-1}$$

$$e_{x, t}(0, 1) = a_{x, t+1}, \quad e_{x, t}(1, 0) = a_{x+1, t} - \theta_1 a_{x+1, t-1} \quad (8.14)$$

$$\hat{z}_{x, t}(1, 1) = -\theta_2 a_{x, t}, \quad e_{x, t}(1, 1) = a_{x+1, t+1} - \theta_1 a_{x+1, t}$$

$$\text{and } \sigma_e^2(0, 1) = \sigma_a^2, \quad \sigma_e^2(1, 0) = \sigma_a^2(1 + \theta_1^2), \quad \sigma_e^2(1, 1) = \sigma_a^2(1 + \theta_1^2).$$

The more general ARMA model is changed to infinite MA model given in (7.5), and terms as far as  $\psi_{10,10}$  carried for accuracy. The method is exactly the same. The models may be updated as new results are obtained.

Forecasts and forecast errors are correlated. In any MA model (and by extension any AR model or ARMA model) the forecast errors are.

$$\begin{aligned} e_{x, t}(0, 1) &= a_{x, t+1}, \quad e_{x, t}(1, 0) = a_{x+1, t} - \theta_1 a_{x+1, t-1}, \\ \text{and} \\ e_{x, t}(1, 1) &= a_{x+1, t+1} - \theta_1 a_{x+1, t}. \end{aligned} \quad (8.15)$$

Hence

$$\begin{aligned} \rho(e_{x, t}(0, 1), e_{x, t}(1, 0)) &= \rho(e_{x, t}(0, 1), e_{x, t}(1, 1)) = 0, \\ \rho(e_{x, t}(1, 0), e_{x, t}(1, 1)) &= -\theta_1(1 + \theta_1^2)^{-1}. \end{aligned} \quad (8.16)$$

Correlations among other forecast errors may be found. Forecasts are also correlated. From (8.14)

$$\rho(\hat{z}_{x, t}(0, 1), \hat{z}_{x, t}(1, 0)) = \rho(\hat{z}_{x, t}(1, 0), \hat{z}_{x, t}(1, 1)) = 0,$$

but

$$\rho(\hat{z}_{x, t}(0, 1), \hat{z}_{x, t}(1, 1)) = \theta_1(\theta_1^2 + \theta_2^2)^{-1/2}. \quad (8.17)$$

The autocorrelation function for forecasts and forecast errors may be found in a similar manner.

It has been assumed in the foregoing that all constants  $\theta_i$  are exact where in fact estimates of  $\theta_i$  are used, including the sample mean  $\bar{z}_{x, t}$ . A discussion of this matter is given in Box and Jenkins (5), pages 267-268. For the mean, an additional source of variation, the  $\sigma_a^2/n$  is added to the formula for  $\sigma_z^2$  given in (8.10) and is important if  $n$  is fairly small,  $n \leq 100$ , depending on the model. For the power spectra of the discussed models, see the papers of Voss et al. (3), Taneja and Aroian (6), and Oprian et al. (7).

#### 9. ACKNOWLEDGMENTS

The suggestions of Robert Perry and Omer Gebizlioglu have been most helpful. Moreover Omer Gebizlioglu has programmed the simulations and performed the many calculations most skillfully. Thanks are due to Dr. Bruno Paul for providing a quiet guest house on Coronado Island where Aroian could work without interruption. This research was supported by the Office of Naval Research, Contract 042-383/78.

#### BIBLIOGRAPHY

- (1) Aroian, L.A. & Gebizlioglu, O. (1980). Time series in m dimensions, spatial models. Unpublished.
- (2) Aroian, L.A. (1979). Multivariate autoregressive time series in m dimensions. Proceedings of the Business and Economics Section, Amer. Stat. Assoc. Annual Meeting, pp. 585-590.
- (3) \*Voss, D., Oprian, C., & Aroian, L.A. (1980). Moving average models, time series in m dimensions.
- (4) Perry, R. & Aroian, L.A. (1979). Of time and the river: time series in m dimensions, the one dimensional autoregressive model. Proceedings of the Statistical Computing Section, Amer. Stat. Assoc. Annual Meeting, pp. 383-388.
- (5) Box, G.E.P. & Jenkins, G.M. (1976). Time Series Analysis: Forecasting and Control, rev. ed. San Francisco: Holden-Day, Inc.
- (6) \*Taneja, V. & Aroian, L.A. (1980). Time Series in m dimensions, autoregressive models.
- (7) \*Oprian, C., Taneja, V., Voss, D., & Aroian, L.A. (1980). General considerations and interrelationships between MA and AR models, time series in m dimensions, the ARMA model.
- (8) \*Aroian, L.A. (1980). Time series in m dimensions.
- (9) Atkinson, A.C. & Pearce, M.C. (1976). The computer generation of beta, gamma, and normal random variables. Journal of the Royal Statistical Society, A, 139, 4, p. 431.

\*Accepted for publication, Commun. Statist., Series B.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AES-8005	2. GOVT ACCESSION NO. A 1733	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Some Simple Examples of Time Series in M Dimensions: An Introduction.	5. TYPE OF REPORT & PERIOD COVERED Technical Report	
6. AUTHOR(s) Leo A. Aroian / Vidya / Taneja	7. PERFORMING ORG. REPORT NUMBER AES-8005	8. CONTRACT OR GRANT NUMBER(s) N00014-77-C-0438
9. PERFORMING ORGANIZATION NAME AND ADDRESS Institute of Administration and Management, Union College & University, Schenectady, NY 12308	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 25 Sep 80	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research, Statistics & Reliabil- ity Program, Office of Naval Research, Arlington, VA 22217	12. REPORT DATE Sept. 25, 1980	13. NUMBER OF PAGES 12
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) Unclassified	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unclassified		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Presented to the 11th Annual Pittsburgh Conference on Modeling & Simulation, May 1-2, 1980, Univ. of Pittsburgh. Paper will appear in the Proceedings of the Conference.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) time series, m dimensions, spatial-time, ARMA models, MA models, AR models, cross correlations, forecasts, forecast errors, power spectrum		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Some simple theoretical examples of time series in m dimensions, their main properties, estimation, and forecasting are given. These are generalizations of the usual time series to events in both time and space. How these models may be simulated is explained. One practical example of flooding on the Mohawk River is given. Included are new results in forecasting; the regions for permissible values of the autocorrelation and cross correlations; and methods for obtaining these correlations.		

DD FORM 1473


JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601Unclassified  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

408725

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)